

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re application of: Morishita et al.

Application No. 09/869,475

Filed: October 10, 2001

Confirmation No. 4309

For: GENE THERAPY FOR DIABETIC
ISCHEMIC DISEASE

Examiner: Brian A. Whiteman

Art Unit: 1635

Attorney Reference No. 6235-59221-01

DECLARATION UNDER § 1.132

1. I, Dr. Ryuichi Morishita, am a co-inventor named in the above-referenced patent application.

2. I have read and understand the above-referenced patent application, including the pending claims, and the Office action dated May 10, 2006.

3. It is my understanding that in the Office action of May 10, 2006, claims 9, 11, 12, 14, 23-26, and 51-52 are rejected under 35 U.S.C. § 103(a) as unpatentable over Isner *et al.* (WO 98/19712), an article from the Japan Financial Times (December 14, 1998), and Li *et al.* (U.S. Patent No. 6,066,123) in further view of Morishita *et al.* (EP 0847757). However, it was not obvious to me and my co-inventor Dr. Toshio Ogihara that diabetic ischemic disease could be treated by administration of hepatocyte growth factor (HGF) gene every few weeks or every few days.

4. At the time the invention was made, it was understood by those skilled in the art such as myself that HGF had a short half-life, such as about 10 minutes. Therefore, it was not expected that the effects of HGF expression would be sustained for a few days or even a few weeks if only administered once over this time period.

5. Several dosages of HGF were tested to identify ranges that permitted the desired therapeutic effect on diabetic ischemia disease. I did not expect that dosages that were therapeutic for arterial disorders would be effective for treating diabetic ischemic disease, as angiogenesis rarely occurs in such diseases.

6. As shown in Exhibit A (Morishita *et al.*, *Hypertension* 44:203-9, 2004), using the claimed methods, two patients (subject numbers 2 and 6 in Table 1) having diabetes mellitus showed therapeutic benefits after administration of HGF gene via an intramuscular injection (Table 2). Two therapeutic injections were administered, four weeks apart. This demonstrates the in vivo effects of administering HGF gene to treat diabetic ischemic disease.

7. I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under 18 U.S.C. § 1001, and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Dr. Ryuichi Morishita

Date

Safety Evaluation of Clinical Gene Therapy Using Hepatocyte Growth Factor to Treat Peripheral Arterial Disease

Ryuichi Morishita, Motokuni Aoki, Naotaka Hashiya, Hirofumi Makino, Keita Yamasaki, Junya Azuma, Yoshiki Sawa, Hikaru Matsuda, Yasufumi Kaneda, Toshio Ogiwara

Abstract—Therapeutic angiogenesis using angiogenic growth factors is expected to be a new treatment for patients with critical limb ischemia (CLI). Because hepatocyte growth factor (HGF) has potent angiogenic activity, we investigated the safety and efficiency of HGF plasmid DNA in patients with CLI as a prospective open-labeled clinical trial. Intramuscular injection of naked HGF plasmid DNA was performed in ischemic limbs of 6 CLI patients with arteriosclerosis obliterans ($n=3$) or Buerger disease ($n=3$) graded as Fontaine III or IV. The primary end points were safety and improvement of ischemic symptoms at 12 weeks after transfection. Severe complications and adverse effects caused by gene transfer were not detected in any patients. Of particular importance, no apparent edema was observed in any patient throughout the trial. In addition, serum HGF concentration was not changed throughout the therapy period in all patients. In contrast, a reduction of pain scale of more than 1 cm in visual analog pain scale was observed in 5 of 6 patients. Increase in ankle pressure index more than 0.1 was observed in 5 of 5 patients. The long diameter of 8 of 11 ischemic ulcers in 4 patients was reduced $>25\%$. Intramuscular injection of naked HGF plasmid is safe, feasible, and can achieve successful improvement of ischemic limbs. Although the present data are conducted to demonstrate the safety as phase I/early phase IIa, the initial clinical outcome with HGF gene transfer seems to indicate usefulness as sole therapy for CLI. (*Hypertension*. 2004;44:203-209.)

Key Words: clinical trials ■ DNA ■ blood vessels ■ safety ■ vascular diseases

The clinical consequences of peripheral arterial disease include pain on walking (claudication), pain at rest, and loss of tissue integrity in the distal ischemic limbs. Although development of beneficial drugs and intervention devices do contribute to the treatment of this disease, critical limb ischemia is estimated to develop in 500 to 1000 individuals per million per year.¹ In a large proportion of these patients, the anatomical extent and the distribution of arterial occlusive disease make the patients unsuitable for operative or percutaneous revascularization. Thus, the disease frequently follows an inexorable downhill course.^{2,3} Recent progress in molecular biology has led to the development of gene therapy⁴⁻¹¹ as a new strategy to treat a variety of cardiovascular diseases. Most of the studies have used vascular endothelial growth factor (VEGF), also known as vascular permeability factor. Indeed, beneficial effects of therapeutic angiogenesis using VEGF gene transfer have been reported in human patients with critical limb ischemia and myocardial ischemia.⁴⁻¹¹ On the other hand, we have focused on hepatocyte growth factor (HGF), because HGF is a potent angio-

genic growth factor in mouse, rat, and rabbit ischemia models.¹²⁻¹⁸ Even in high-risk conditions for atherosclerosis such as a diabetic and high lipoprotein (a) concentration model, overexpression of HGF is enough to stimulate collateral formation to treat ischemic symptoms.^{17,18} Unexpectedly, the mitogenic activity of HGF has been reported to be more potent than that of VEGF in human aortic endothelial cells in vitro as well as a rabbit hindlimb ischemia model in vivo.^{13,19} Based on our previous data, we designed a human clinical trial of gene therapy for peripheral artery disease using HGF gene as an open-labeled study. Despite this limitation of the trial, we demonstrate that intramuscular injection of naked HGF plasmid is safe and feasible as subanalysis of phase I/IIa.

Methods

Plasmid DNA

The pVAX1 plasmid vector (3.0 kb) is commercially produced by Invitrogen Corporation (Carlsbad, Calif) and was selected for HGF construct because it incorporated elements to be in compliance with

Received January 9, 2004; first decision January 23, 2004; revision accepted June 9, 2004.

From the Department of Geriatric Medicine (R.M., M.A., N.H., H.M., K.Y., J.A., T.O.), Division of Clinical Gene Therapy (R.M.), Department of Surgery (Y.S., H.M.), and Division of Gene Therapy Science (Y.K.), Graduate School of Medicine, Osaka University, Japan.

R.M. is a board member and stockholder of AnGes MG, Inc, which has developed HGF gene as a gene therapy drug. T.O., M.A., N.H., H.M., K.Y., J.A., and Y.K. are also stockholders of AnGes MG, Inc.

Correspondence to Ryuichi Morishita, MD, PhD, Professor, Division of Clinical Gene Therapy, Graduate School of Medicine, Osaka University, 2-2 Yamada-oka, Suita 565-0871, Japan. E-mail morishit@cg.t-med.osaka-u.ac.jp

© 2004 American Heart Association, Inc.

Hypertension is available at <http://www.hypertensionaha.org>

DOI: 10.1161/01.HYP.0000136394.08900.ed




TABLE 1. Patient Characteristics

No.	Age (Sex)	Disease	Fontaine	Rutherford Classification	Risk Factors Classification	Previous for Arteriosclerosis	DSA/MRA Treatment
1	57 (M)	ASO	III	II-4	Smoking, HT	7 Bypass grafts (occluded) 2PTA Sympathectomy Prostaglandin therapy Antiplatelet drugs (clostazol, ticlopidine, beraprost)	SFA, POP, PT, PN occlusion
2	42 (M)	ASO	IV	III-5	Smoking, HT, DM CRF (HD)	Sympathectomy Prostaglandin therapy Antiplatelet drugs (clostazol, ticlopidine, beraprost)	SFA, POP, AT occlusion
3	48 (M)	Buerger	IV	III-5	—	Bypass graft (occluded) Prostaglandin therapy Antiplatelet drugs (clostazol, ticlopidine, beraprost)	SFA, POP, AT, PT, PN occlusion
4	64 (F)	Buerger	III	II-4	—	Prostaglandin therapy Antiplatelet drugs (ticlopidine, beraprost)	SFA, POP, AT, PT, PN occlusion
5	67 (M)	Buerger	IV	III-5	—	Prostaglandin therapy Antiplatelet drugs (ticlopidine)	AT, PT, PN occlusion
6	69 (M)	ASO	IV	III-5	HT, DM	Bypass graft (occluded) Prostaglandin therapy Antiplatelet drugs (clostazol, ticlopidine, beraprost)	POP, AT, PN occlusion

HT indicates hypertension; DM, diabetes mellitus; CRF, chronic renal failure; HD, hemodialysis; SFA, superficial femoral artery; POP, popliteal; AT, anterior tibial artery; PT, posterior tibial artery; PN, peroneal artery.

the Food and Drug Administration document, *Points to Consider on Plasmid DNA Vaccines for Preventative Infectious Disease Indications*. To minimize the possibility of chromosomal integration, insertional mutagenesis through the activation of oncogenes, or inactivation of tumor suppressor genes, all sequences with possible DNA, along with any sequences not necessary for replication in *Escherichia coli* or for expression of recombinant protein in mammalian cells. pVAX1HGF plasmid consists of the cDNA fragment of human HGF (*EcoRI*/NotI, 2.2 kbp) inserted into the pVAX1 vector. The construct does not have DNA sequence homology to the human genome (except the HGF insert), and the vector sequences do not contain known oncogenic potential.

Patient Cohort

In this initial single-center phase of a prospective open-labeled study, patients could be enrolled if they (1) had chronic critical limb ischemia, including rest pain or nonhealing ischemic ulcers, for a minimum 4 weeks; (2) were resistant to conventional drug therapy at least for >4 weeks after hospitalization; (3) were not candidates for surgical or percutaneous revascularization based on usual practice standards; (4) did not have cancer or a history of cancer; and (5) did not have severe unstable retinopathy. Objective documentation of ischemia, including a resting ankle brachial index (ABI) of less than 0.6 in the affected limb on 2 consecutive examinations performed 1 week apart was necessary. Patients were observed for 4 weeks under conventional drug therapy to confirm that their clinical symptoms and objective parameters were not improved. Selection criteria was confirmed by an independent committee for assessment and evaluation of clinical gene therapy at Osaka University, which was approved by the Ministry of Welfare and the Ministry of Education (Science and Culture). After confirmation, 6 patients (5 male and 1 female, aged 57.8±4.5 years) with arteriosclerosis obliterans (ASO; n=3) or Buerger disease (n=3) graded as Fontaine III or IV underwent direct intramuscular gene transfer of naked plasmid DNA encoding HGF. The characteristics of the patients are shown in Table 1.

Intramuscular Injection of Naked Plasmid DNA Encoding Human HGF

Each patient received an intramuscular injection of naked plasmid DNA encoding HGF gene regulated by the cytomegalovirus promoter/enhancer. The preparation and purification of the plasmid from

cultures of HGF-transformed *E. coli* was performed at Qiagen, Inc using kanamycin. The purified plasmid was transported to the human gene therapy laboratory at Osaka University Medical School and stored in vials at -20°C. Some of them were pooled for quality-control analysis and sequence check. No vectors were used to deliver plasmid.

First, test intramuscular injection of small dose (test injection; 0.4 mg plasmid DNA) was performed to examine acute or subacute allergy to plasmid DNA (Table 1, available online at <http://www.hypertensionaha.org>). After confirmation of no allergic reaction or anaphylaxis, a therapeutic dose (2 mg) of naked HGF plasmid DNA was intramuscularly injected 2 weeks after test injection. Five hundred micrograms of HGF plasmid DNA was diluted in a sterile saline solution up to 3 mL, and 4 aliquots (total 2000 µg/12 mL) were administered into the calf or distal thigh muscles of the ischemic limbs by direct intramuscular injection under echonographic guidance. Four injection sites were selected arbitrarily, according to the angiographic findings and the available muscle mass. The injection sites of plasmid DNA are summarized in Figure 1. Four weeks after the initial injection, a second injection (2000 µg) was similarly administered, giving a total dose of 4000 mg plasmid DNA per patient.

Patient Follow-Up and Assessment

Patients were followed-up by physical examination (including change in ischemic ulcers), blood analysis, visual analog scale (pain scale), SF-36 (quality-of-life scale), transcutaneous PO₂ (TcPO₂), and measurements of ABI and toe pressure index (TPI) weekly during the first 12 weeks, every other week for the next 8 weeks, monthly for the next 12 weeks, and every 3 months up to 2 years after the first injection of plasmid DNA (Table 1). All patients received usual and full drug therapy for peripheral arterial disease including antiplatelet drugs, which were not changed throughout this trial. Ischemic ulcers and necrotic lesions were managed in a standard fashion, and antibiotics were prescribed as necessary. ABI and TPI were measured using the toe plethysmographic wave or Doppler wave with an IMEX (GETS Brothers). TcPO₂ was calculated before and after supply of O₂ (5 L/min×5 minutes) (Sumitomo-denno Hitechs Inc). Intra-arterial digital subtraction angiography, magnetic resonance angiography, and, if possible, computerized tomographic angiography were performed within 1 month before the first injection and 4 weeks after each treatment and 3 months after




Figure 2. Angiogram in patient No. 3. DSA from patient No. 3 with Burger disease. An arrow indicates a newly recanalized vessel.

was shown in 2 of 6 treated ischemic limbs. Figure 2 demonstrates DSA findings in a patient with Burger disease (No. 3). A large vessel, which is indicated by an arrow, was newly observed (Figure 2a to 2c). Although it is not clear whether this vessel was new or not, it is possible that an increase of new microvessels led to the recanalization. This recanalization was also confirmed by serial magnetic resonance angiograms (data not shown). DSA of another patient with Burger disease (No. 5) showed a marked increase in peripheral blood flow and formation of new blood vessels (Figure 3a and 3b).

To evaluate the functional improvement of HGF gene therapy, we also measured ABI during gene therapy. Although ABI could not be measured in 1 patient because of




Figure 3. Angiogram in patient No. 5. DSA from patient No. 5 with Burger disease. Peripheral blood flow is markedly increased.




Figure 4. Change in ABI in 5 limbs after intramuscular injection of HGF plasmid DNA. Baseline indicates 4 weeks before test injection; Pre, just before test injection; 4 weeks, 4 weeks after first injection (just before second injection); 8 weeks, 8 weeks after first injection (4 weeks after second injection); 12 weeks, 12 weeks after first injection (8 weeks after second injection). * $P < 0.05$ vs baseline.

uncompressible severely calcified vessels, ABI was significantly increased from 0.426 ± 0.046 ($n = 5$) at baseline (before administration) to 0.626 ± 0.071 ($P = 0.0155$; $n = 5$) at 4 weeks after the second injection and to 0.596 ± 0.046 ($P = 0.0360$; $n = 5$) at 8 weeks after the second injection (Figure 4). The absolute value of systolic ankle pressure was significantly increased in 5 limbs after gene transfer, whereas ankle pressures of untreated limbs were not significantly changed (data not shown). Also, TPI, which could be measured only in 2 patients (Nos. 1 and 3), tend to be increased (data not shown), accompanied by an improvement of ABI. However, TPI was not measured in 4 patients (No. 2, 4, 5, and 6) because of ischemic ulcers on the great toes of their ischemic legs. When an increase in ABI of >0.1 was assumed to be an improvement, according to the standard of Rutherford, 5 of 5 patients (100%) showed a positive response. In addition, as transcutaneous PO_2 is an indicator of the effectiveness in terms of angiogenesis and increase in blood supply in targeted ischemic lesions, we also measured $TcPO_2$. As shown in Table 3, the change in $TcPO_2$ after O_2 stimulation was significantly increased at 8 weeks compared with baseline ($P < 0.05$).

To evaluate the effects of HGF gene therapy in clinical symptoms, we used the change in the ischemic ulcer and visual analogue scale. In this trial, a total of 11 ischemic ulcers were found in 4 patients. As shown in Figure 5, 2 of 11 ulcers completely disappeared. Considering an improvement of ischemic ulcers of more than 25% to be evaluated as positive, 8 of 11 ulcers (72%) improved. Typical examples of

TABLE 3. Transition of $TcPO_2$

Week	0 Week	4 Weeks	8 Weeks	12 Weeks
Pre O_2	58.7 ± 2.8	60.2 ± 5.9	45.8 ± 11.3	68.3 ± 8.0
Post O_2	80.3 ± 6.7	83.2 ± 5.4	69.3 ± 17.4	$99.5 \pm 13.4^*$
Δ (Post-pre)	21.7 ± 4.5	23.0 ± 3.7	23.5 ± 9.0	$31.2 \pm 6.4^*$

* $P < 0.05$ vs 0 week.




Figure 5. Change in size of ischemic ulcers in 4 limbs after intramuscular injection of HGF plasmid DNA. Baseline indicates 4 weeks before test injection; Pre, just before test injection; 4 weeks, 4 weeks after first injection (just before second injection); 8 weeks, 8 weeks after first injection (4 weeks after second injection); 12 weeks, 12 weeks after first injection (8 weeks after second injection). * $P < 0.05$ vs baseline.

the change in ischemic ulcers in patients with Buerger disease (No. 3 and 5) are shown in Figure 6. Three of 4 patients demonstrated an improvement of the maximum ischemic ulcer diameter of $>25\%$ (efficacy rate = 75%). Also, we evaluated resting pain using a visual analog scale, as a standard method for the evaluation of pain, where 0.0 cm means "pain free" or no pain, and 10 cm means more severe pain. As shown in Figure 7, pain was significantly improved in a time-dependent manner.

Discussion

The natural history of critical limb ischemia has been well documented to have an inexorable downhill course.¹ The inclusion criteria for this study were drafted to restrict treatment to patients in whom the natural history of critical limb ischemia had been established previously. Especially, this trial excluded patients who demonstrated an improvement of clinical symptoms after hospitalization and pharmacological treatment, to identify the effects of HGF gene




Figure 6. Typical examples of change in ischemic ulcers in patients with Buerger disease (patient Nos. 3 and 5). Pre-GT indicates 4 weeks before test injection; 8 weeks, 8 weeks after first injection (4 weeks after second injection).




Figure 7. Reduction of visual analog scale after intramuscular injection of HGF plasmid DNA. Baseline indicates 4 weeks before test injection; Pre, just before test injection; 4 weeks, 4 weeks after first injection (just before second injection); 8 weeks, 8 weeks after first injection (4 weeks after second injection); 12 weeks, 12 weeks after first injection (8 weeks after second injection). * $P < 0.05$ vs baseline.

therapy on clinical symptoms. In addition, the evaluation of efficacy and safety was performed by an independent committee without the researchers who carried out this trial. Four of the 6 limbs had developed frank gangrene. Although inclusion criteria required a minimum of 4 weeks of conservative measures without evidence of improvement, in reality, signs or symptoms of critical limb ischemia had been present in all cases for 2 to 10 months before gene therapy. In this series of 6 patients, 6 developed critical limb ischemia despite having undergone many vascular surgical reconstructions. Furthermore, because HGF had not been administered previously as recombinant protein or gene therapy, no data were available from any source to indicate either the safety or bioactivity of any dose of plasmid human HGF DNA. Accordingly, the design of this phase I/early phase II trial, approved by the Japanese Government, was conducted as a nonrandomized consecutive treatment series based on the animal experiments.^{15, 18, 22, 24} The previous articles reported the detection of human HGF protein in mouse experiments (control; not detected, HGF: 1.05 ± 0.08 ng/g tissue¹⁵), rat experiments (control; not detected, HGF: 1.95 ± 0.42 ng/g tissue¹⁸), and rabbit experiments (control; not detected, HGF: 2.08 ± 0.39 ng/g tissue^{15, 24}).

In all patients, treatment was shown to achieve clinically significant modulation of the recipient phenotype. Noninvasive studies documented hemodynamic evidence of improved limb perfusion that satisfied the outcome criteria proposed to assess the results of surgical reconstruction or percutaneous revascularization.²⁵ Absolute ankle pressure increased in 5 limbs after gene therapy. ABI increased from 0.426 ± 0.046 at baseline to 0.596 ± 0.046 at 12 weeks after the first injection. To put this in perspective, an increase in ABI of >0.1 is considered indicative of a successful surgical or percutaneous intervention.²⁵ This increase in ABI was similar to the results of VEGF trials.^{5, 6} To our knowledge, such an improvement has not previously been achieved spontaneously or with medical therapy in patients with critical limb ischemia.

Similarly, angiographic demonstration of newly visible collateral vessels, accompanied here by noninvasive evidence of improved blood flow, confirmed these improvements, although the quantification of angiograms is quite difficult. Indeed, previous reports have indicated that current methods used to perform diagnostic contrast angiography cannot provide images of arteries measuring $< 200 \mu\text{m}$ in diameter.²⁶ The conventional angiographic techniques used in this study may have failed to depict the full extent of angiogenesis achieved after HGF gene transfer. Rest pain was significantly improved from baseline (before treatment) to 12 weeks after transfection as assessed by visual analog scale. Ischemic ulcers healed or improved markedly by $>25\%$ in 3 of 4 patients; this included 3 patients recommended for below-knee amputation in whom successful limb salvage was achieved. Given the poor prognosis for patients with chronic critical limb ischemia, in whom the possibility of spontaneous improvement is remote,²⁷ the outcome in this initial cohort is thus encouraging. Nevertheless, the present study cannot fully eliminate the placebo effects. Further studies in a randomized placebo-double blinded manner are necessary to demonstrate the effectiveness of HGF gene therapy.

Unlike previous reports using VEGF gene showing that VEGF protein level demonstrated a transient peak in the systemic circulation 1 to 3 weeks after gene transfer, our trial did not demonstrate an increase in serum HGF concentration during gene therapy. It is noteworthy that there was no evidence of edema in the patients transfected with the human HGF gene, in marked contrast to a VEGF trial in which 60% of patients developed moderate or severe edema in a phase I/IIa trial. What is the difference between HGF and VEGF? We believe that one of the distinguishing features of HGF is that it stimulates the migration of vascular smooth muscle cells (VSMCs) without the replication of VSMCs, whereas VEGF does not stimulate either the migration or proliferation of VSMCs because of its lack of receptors in VSMCs. The initial event in angiogenesis induced by VEGF is the migration of endothelial cells, leading to the sprouting of blood vessels. Later, the migration of VSMCs occurs because of the release of platelet-derived growth factor, followed by the migration of endothelial cells. However, a delay in the maturation of blood vessels might exist in the case of angiogenesis induced by VEGF. In contrast, HGF simultaneously stimulated the migration of both endothelial cells and VSMCs. Thus, the blood vessels may mature at an earlier time point, thereby avoiding the release of blood-derived cells into the extracellular space, although further studies might be necessary to examine the angiogenic properties of various angiogenic growth factors including HGF, VEGF, and fibroblast growth factor.

Regarding the safety of HGF gene therapy, previous work established that transgene expression using plasmid DNA is limited to <30 days in animal models of limb ischemia.^{27,28} In contrast, it appears that in both animals and humans, collateral vessel development sufficient to restore limb perfusion to satisfactory resting levels occurs within this time interval. The cessation of gene expression beyond this time point can be considered to constitute an inherent safety feature of HGF gene transfer that protects the organism from

indefinite constitutive expression of an angiogenic growth factor. In addition, the circulating level of HGF is elevated in patients with hypertension, peripheral arterial disease, and myocardial infarction,^{21,29,30} although the present study demonstrated no increase in serum HGF concentration during gene therapy. Potential side effects, such as cancer or diabetic retinopathy due to an increase in plasma HGF secreted from transfected cells, would be minimized in gene therapy using intramuscular injection of HGF into ischemic muscle. Nevertheless, these findings are preliminary and do not establish the long-term safety of HGF. Clearly, further clinical studies of alternative dosing regimens of gene therapy with randomized placebo-controlled trials will be required to define the efficacy of this therapy.

Acknowledgments

This work was partially supported by grants-in-aid from the Organization for Pharmaceutical Safety and Research, the Ministry of Public Health and Welfare, and Japan Promotion of Science, and through special coordination funds of the Ministry of Education, Culture, Sports, Science and Technology, the Japanese Government.

References

- Second European Consensus Document on Chronic Critical Leg Ischemia. *Circulation*. 1991;84(Suppl IV):IV-1-IV-26.
- Dormandy J, Mahir M, Ascaly G, Balsano F, De-Leuze P, Blombery P, Bousset MG, Clement D, Coffman J, Deuschle M, Alety O, Hampton J, Hahler E, Ohlin P, Rieger H, Strandén E, Turpie AGG, Uriai L, Verstraete M. Fate of the patient with chronic leg ischaemia. A review article. *J Cardiovasc Surg (Torino)*. 1989;30:50-57.
- Rutherford RB, Flanigan DP, Gupta SK, Johnston KW, Karmody A, Whittemore AD, Baker D, Ernst CB, Jamieson C, Mehta S. Suggested standards for reports dealing with lower extremity ischemia. Ad Hoc Committee on Reporting Standards. Society for Vascular Surgery/North Am Chapter, International Society for Cardiovascular Surgery. *J Vasc Surg*. 1986;4:80-94.
- Inzer JM, Baumgartner I, Rauh G, Schainfeld R, Blair R, Manor O, Razvi S, Symes JF. Treatment of thromboangiitis obliterans (Buerger's disease) by intramuscular gene transfer of vascular endothelial growth factor: preliminary clinical results. *J Vasc Surg*. 1998;28:964-973.
- Baumgartner I, Pieczek A, Manor O, Blair R, Kearney M, Walsh K, Inzer JM. Constitutive expression of phVEGF165 after intramuscular gene transfer promotes collateral vessel development in patients with critical limb ischemia. *Circulation*. 1998;97:1114-1123.
- Baumgartner I, Rauh G, Pieczek A, Wanssch D, Magner M, Kearney M, Schainfeld R, Inzer JM. Lower-extremity edema associated with gene transfer of naked DNA encoding vascular endothelial growth factor. *Ann Intern Med*. 2000;132:880-884.
- Losordo DW, Vale PR, Symes JF, Dunnington CH, Esakof DD, Mayasky M, Ashare AB, Lathi K, Inzer JM. Gene therapy for myocardial angiogenesis: initial clinical results with direct myocardial injection of phVEGF165 as sole therapy for myocardial ischemia. *Circulation*. 1998;98:2800-2804.
- Vale PR, Losordo DW, Milliken CE, Esakof DD, Inzer JM. Images in cardiovascular medicine: Percutaneous myocardial gene transfer of phVEGF-2. *Circulation*. 1999;100:2462-2463.
- Vale PR, Losordo DW, Milliken CE, Mayasky M, Esakof DD, Symes JF, Inzer JM. Left ventricular electroanatomic mapping to assess efficacy of phVEGF165 gene transfer for therapeutic angiogenesis in chronic myocardial ischemia. *Circulation*. 2000;102:965-974.
- Rosengart TK, Lee LY, Patel SR, Sanborn TA, Parikh M, Bergman GW, Hachamovitch R, Szulc M, Kilgfield PD, Okin PM, Hahn RT, Devereux RB, Post MR, Hackett NR, Foster T, Grasso TM, Lesser ML, Isom OW, Crystal RG. Angiogenesis gene therapy: phase 1 assessment of direct intramyocardial administration of an adenoviral vector expressing VEGF121 cDNA to individuals with clinically significant severe coronary artery disease. *Circulation*. 1999;100:468-474.
- Rosengart TK, Lee LY, Patel SR, Kilgfield PD, Okin PM, Hackett NR, Isom OW, Crystal RG. Six-month assessment of a phase 1 trial of angiogenic gene therapy for the treatment of coronary artery disease using

- direct intramyocardial administration of an adenovirus vector expressing the VEGF121 cDNA. *Ann Surg*. 1999;230:466-470.
12. Morishita R, Nakamura S, Hayashi S, Taniyama Y, Moriguchi A, Nagano T, Tajiri M, Noguchi H, Takeshita S, Matsumoto K, Nakamura T, Higaki J, Ogihara T. Therapeutic angiogenesis induced by human recombinant hepatocyte growth factor in rabbit hind limb ischemia model as cytokine supplement therapy. *Hypertension*. 1999;33:1379-1384.
 13. Belle EV, Witzenschnitzer B, Chen D, Silver M, Chang L, Schwall R, Inzer JM. Potentiated angiogenic effect of scatter factor/hepatocyte growth factor by induction of vascular endothelial growth factor: the case for paracrine amplification of angiogenesis. *Circulation*. 1998;97:381-390.
 14. Hayashi S, Morishita R, Nakamura S, Yamamoto K, Moriguchi A, Nagano T, Tajiri M, Noguchi H, Matsumoto K, Nakamura T, Higaki J, Ogihara T. Potential role of hepatocyte growth factor, a novel angiogenic growth factor, in peripheral arterial disease: down-regulation of HGF in response to hypoxia in vascular cells. *Circulation*. 1999;100:11301-11308.
 15. Taniyama Y, Morishita R, Aoki M, Nakagami H, Yamamoto K, Yamazaki K, Matsumoto K, Nakamura T, Kaneda Y, Ogihara T. Therapeutic angiogenesis induced by human hepatocyte growth factor gene in rat and rabbit hind limb ischemia models: preclinical study for treatment of peripheral arterial disease. *Gene Ther*. 2001;8:181-189.
 16. Aoki M, Morishita R, Taniyama Y, Kida I, Moriguchi A, Matsumoto K, Nakamura T, Kaneda Y, Higaki J, Ogihara T. Angiogenesis induced by hepatocyte growth factor in non-infarcted myocardium and infarcted myocardium: up-regulation of essential transcription factor for angiogenesis, ets. *Gene Ther*. 2000;7:417-427.
 17. Taniyama Y, Morishita R, Hirooka K, Aoki M, Nakagami H, Yamazaki K, Matsumoto K, Nakamura T, Kaneda Y, Ogihara T. Therapeutic angiogenesis induced by human hepatocyte growth factor gene in rat diabetic hind limb ischemia model: molecular mechanisms of delayed angiogenesis in diabetes. *Circulation*. 2001;104:2344-2350.
 18. Morishita R, Sakaki M, Yamamoto K, Iguchi S, Aoki M, Yamazaki K, Matsumoto K, Nakamura T, Lawn R, Ogihara T, Kaneda Y. Impairment of collateral formation in I(p6) transgenic mice: therapeutic angiogenesis induced by human hepatocyte growth factor gene. *Circulation*. 2002;105:1491-1496.
 19. Nakamura Y, Morishita R, Higaki J, Kida I, Aoki M, Moriguchi A, Yamada K, Hayashi S, Yo Y, Nakano H, Matsumoto K, Nakamura T, Ogihara T. Hepatocyte growth factor is a novel member of the endothelium-specific growth factors: additive stimulatory effect of hepatocyte growth factor with basic fibroblast growth factor but not with vascular endothelial growth factor. *J Hypertens*. 1996;14:1067-1072.
 20. Yamada A, Matsumoto K, Iwanami H, Sekiguchi K, Kawata S, Matsuzawa Y, Nakamura T. Rapid and sensitive enzyme-linked immunosorbent assay for measurement of HGF in rat and human tissues. *Biomed Res*. 1995;16:105-114.
 21. Nakamura Y, Morishita R, Nakamura S, Aoki M, Moriguchi A, Matsumoto K, Nakamura T, Higaki J, Ogihara T. A vascular modulator, hepatocyte growth factor, is associated with systolic pressure. *Hypertension*. 1996;28:409-413.
 22. Tomita N, Morishita R, Taniyama Y, Koike H, Aoki M, Shimizu H, Matsumoto K, Nakamura T, Kaneda Y, Ogihara T. Angiogenic property of hepatocyte growth factor is dependent on up-regulation of essential transcription factor for angiogenesis, ets-1. *Circulation*. 2003;107:1411-1417.
 23. Hirooka K, Koike H, Yamamoto S, Tomita N, Yokoyama C, Tanabe T, Aikou T, Ogihara T, Kaneda Y, Morishita R. Enhanced therapeutic angiogenesis by co-transfection of prostacyclin synthase gene or optimization of intramuscular injection of naked plasmid DNA. *Circulation*. 2003;108:2689-2696.
 24. Koike H, Morishita R, Iguchi S, Aoki M, Matsumoto K, Nakamura T, Yokoyama C, Tanabe T, Ogihara T, Kaneda Y. Enhanced angiogenesis and improvement of neuropathy by co-transfection of human hepatocyte growth factor and prostacyclin synthase gene. *FASEB J*. 2003;17:779-781.
 25. Rutherford RB, Becker GJ. Standards for evaluating and reporting the results of surgical and percutaneous therapy for peripheral arterial disease. *Radiology*. 1991;181:277-281.
 26. Takeshita S, Ishiki T, Tanaka E, Eto K, Miyazawa Y, Tanaka A, Shinzaki Y, Hyodo K, Ando M, Kubota M, Tanaka K, Umetsu K, Ochiai M, Sato T, Mori H, Miyashita H. Use of synchrotron radiation microangiography to assess development of small collateral arteries in a rat model of hindlimb ischemia. *Circulation*. 1997;95:805-808.
 27. Takeshita S, Ishiki T, Sato T. Increased expression of direct gene transfer into skeletal muscles observed after acute ischemic injury in rats. *Lab Invest*. 1996;74:1061-1065.
 28. Tsurumi Y, Takeshita S, Chen D, Kearney M, Rossow ST, Passeri J, Horowitz JR, Symes JF. Direct intramuscular gene transfer of naked DNA encoding vascular endothelial growth factor augments collateral development and tissue perfusion. *Circulation*. 1996;94:3281-3290.
 29. Yoshitomi Y, Kojima S, Umemoto T, Kubo K, Matsumoto Y, Yano M, Sugi T, Kuramochi M. Serum hepatocyte growth factor in patients with peripheral arterial occlusive disease. *J Clin Endocrinol Metab*. 1999;84:2425-2458.
 30. Nakamura S, Moriguchi A, Morishita R, Aoki M, Yo Y, Hayashi S, Nakano N, Katsuya T, Nakata S, Takami S, Matsumoto K, Nakamura T, Higaki J, Ogihara T. A novel vascular modulator, hepatocyte growth factor (HGF), as a potential index of the severity of hypertension. *Biochem Biophys Res Commun*. 1998;242:238-243.